

Date: March 2019













Skin and Wound Product Information Sheet

Cavilon Advanced Skin Protectant		
Classification	Advanced Skin Protectant: Film Barri	er
Key Points	moist/wet non-intact skin.	crylate solution used to protect dry intact and roof, durable, flexible, breathable, transparent film
	or caustic irritants. • Product wears off of the skin over time; do	
Indications	as per, NSWOC/Wound Clinician when othe the skin from: • worsening Moisture Associated Skin D o on perineal/buttock skin(Inconti o on peri-wound, peri-ostomy, per o in skin folds • skin damage related to a dhesive dress	ultation with, and after receiving application instruction r skin protectants have not been effective in protecting ramage(MASD): nent Associated Dermatitis (IAD))
Precautions	Skin Injury (MARSI)) • Extremely flammable until it has completely dried on the skin, do not apply in the presence of fire ignition or heat-producing sources; use in a well-ventilated space • The product has not been tested for use on babies/infants/neonates	
	or other materials (tapes, dressings, ostom • May enhance the adhesiveness of adhesiveness of a dhesiveness	ning into contact with other skin a reas (e.g. skin fold) by appliances) e products; e.g. tape, ostomy barrier, especially within MARSI, use an adhesive remover when removing such
Contraindications	 Do not apply to full-thickness wounds Do not apply to second or third degree bur Do not apply to infected areas 	ches or with medicated powders, ointments or creams faround the eyes
Formats & Sizes	2.7ml Applicator (covers an area of approximately 25.4x 25.4cm)	
1	Application Directions	Rationale
For the initial application, cleanse the skin with appropriate cleanser for the area; e.g. a no-rinse skin cleanser, water, normal saline; ensuring all trace of previous skin care product is removed (moisturizer, silicone or zinc protectant, petro-latum). Gently pat the skin dry.		Solution does not a dhere to skin that has a moisturizer / barrier product on it.
For denuded/eroded skin, gently pat the surface with gauze to blot any excessive moisture.		Solution does not adhere to skin that is excessively moist.
Grasp the applicator, place thumb at the end of the lever, aim the applicator down and then tilt the sponge tip <u>down</u> as well.		Applicator needs to be held in a downward position to allow the solution to flow into the sponge Ensuring that the sponge tip is tilted down will allow the solution to flow into the tip.



Date: March 2019













Skin and Wound Product Information Sheet

Skin and Wound Product	information Sheet
Firmly depress the lever to break the internal ampule. A snapping/popping sound will be heard when the ampule is	
broken. Release the thumb pressure on the lever.	
Continue to hold the applicator in a downward/tilted position for	
at least 10 seconds to allow the solution to flow into the tip of	
foam sponge. Solution will not completely saturate the foam	
sponge to the side edges. Use one hand to hold the skin surface taut/lift up the skin fold and	Holding the chin surface to ut will a sciet with
apply the solution using a gentle, even, <u>continuous</u> sweeping	Holding the skin surface taut will assist with minimizings kin wrinkles or folds making it easier to
(back &forth) motion over the area of concern with minimal overlapping.	apply the solution in a continuous sweeping motion.
Do not apply pressure to the lever or on the sponge as this may	
result in excessive solution in the sponge causing the area to be	
too wet.	
Allow area to thoroughly dry for at least 60 seconds before it	Allowing a dequate dry-time will avoid unintentional skin a dherence.
come into contact with other skin areas (e.g. skin folds), adhesive	skin adherence.
products (tape, bordered dressing, ostomy appliance), clothing, incontinence products, etc.	
If an area has been missed, wait the 60 seconds before applying the solution to that area.	
Frequency of Re-application	
Inspect the area at least every shift; cleanse the area as needed.	If applied too frequently will cause excess build-up.
Reapply another thin layer every 3-4 days.	
Discontinue use of the product when skin is healthy or 'step-	
down' to a nother skin care intervention to continue the healing	
1 down to another skin care miter vention to continue the meaning	
process.	
_	
process. To Remove Film will naturally wear off. If the film must be removed, then use	Product wears off as epidermal cells naturally slough
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethy-	Product wears off as epidermal cells naturally slough off.
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethy-disiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for	•
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethydisiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin.	•
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydis iloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes,	•
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydis iloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied	· · · · · · · · · · · · · · · · · · ·
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethydisiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier.	
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydis iloxane (HMDS) e.g (3M Adhes ive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations	off.
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydisiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and	· · · · · · · · · · · · · · · · · · ·
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydis iloxane (HMDS) e.g (3M Adhes ive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations	Approximately 2% of the population may have an
To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydis iloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an	Approximately 2% of the population may have an
process. To Remove Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethydis iloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water.	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethy-disiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention.	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydisiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethydisiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary.	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethydisiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary. If skin or other materials become stuck to the area, use an	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containings ilicones or hexamethydisiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary. If skin or other materials become stuck to the area, use an adhesive remover appropriate for client's skin to separate the two	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethy-disiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary. If skin or other materials become stuck to the area, use an adhesive remover appropriate for client's skin to separate the two skin surfaces or to separate the material from the barrier.	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethy-disiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary. If skin or other materials become stuck to the area, use an adhesive remover appropriate for client's skin to separate the two skin surfaces or to separate the material from the barrier. Expected Outcome	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethy-disiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary. If skin or other materials become stuck to the area, use an adhesive remover appropriate for client's skin to separate the two skin surfaces or to separate the material from the barrier.	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containingsilicones or hexamethy-disiloxane (HMDS) e.g (3 M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary. If skin or other materials become stuck to the area, use an adhesive remover appropriate for client's skin to separate the two skin surfaces or to separate the material from the barrier. Expected Outcome If used for the management of MASD, film barrier prevented further MASD.	Approximately 2% of the population may have an
Film will naturally wear off. If the film must be removed, then use an adhesive remover containing silicones or hexamethy-disiloxane (HMDS) e.g (3M Adhesive Remover) appropriate for client's skin. Use an adhesive remover when removing products such as tapes, bordered dressings, ostomy appliances that have been applied over top of the film barrier. To Manage Adverse Situations If irritation or pain develops (redness, inflammation, heat, and change in skin colour) discontinue product use and use an adhesive remover containing silicones or HMDS to remove as much of the film barrier as possible. If solution enters the eye, immediately flush with lukewarm water. Do not rub or try to separate the eyelids. Seek medical attention. If there is an accidental contact with a wound with depth, flush with water or saline; complete removal is not necessary. If skin or other materials become stuck to the area, use an adhesive remover appropriate for client's skin to separate the two skin surfaces or to separate the material from the barrier. Expected Outcome If used for the management of MASD, film barrier prevented	Approximately 2% of the population may have an